

2007

USP 30

THE UNITED STATES PHARMACOPEIA

NF 25

THE NATIONAL FORMULARY

Volume 1

*By authority of The United States Pharmacopeial Convention, meeting at Washington, D.C., March 9–13, 2005. Prepared by the Council of Experts and published by the Board of Trustees*

*Official from May 1, 2007*

The designation on the cover of this publication, “USP NF 2007,” is for ease of identification only. The publication contains two separate compendia: *The United States Pharmacopeia*, Thirtieth Revision, and the *National Formulary*, Twenty-Fifth Edition.

THE UNITED STATES PHARMACOPEIAL CONVENTION  
12601 Twinbrook Parkway, Rockville, MD 20852

Where drying in vacuum over a desiccant is directed in the individual monograph, a vacuum desiccator or a vacuum drying pistol, or other suitable vacuum drying apparatus, is to be used.

**Water**—Where water is called for in tests and assays, *Purified Water* is to be used unless otherwise specified. For special kinds of water such as "carbon dioxide-free water," see the introduction to the section *Reagents, Indicators, and Solutions*. For *High-Purity Water* see *Containers* (661).

**Water and Loss on Drying**—Where the water of hydration or adsorbed water of a Pharmacopeial article is determined by the titrimetric method, the test is generally given under the heading *Water*. Monograph limits expressed as a percentage are figured on a weight/weight basis unless otherwise specified. Where the determination is made by drying under specified conditions, the test is generally given under the heading *Loss on drying*. However, *Loss on drying* is most often given as the heading where the loss in weight is known to represent residual volatile constituents, including organic solvents as well as water.

**Test Results, Statistics, and Standards**—Interpretation of results from official tests and assays requires an understanding of the nature and style of compendial standards, in addition to an understanding of the scientific and mathematical aspects of laboratory analysis and quality assurance for analytical laboratories.

Confusion of compendial standards with release tests and with statistical sampling plans occasionally occurs. Compendial standards define what is an acceptable article and give test procedures that demonstrate that the article is in compliance. These standards apply at any time in the life of the article from production to consumption. The manufacturer's release specifications, and compliance with good manufacturing practices generally, are developed and followed to ensure that the article will indeed comply with compendial standards until its expiration date, when stored as directed. Thus, when tested from the viewpoint of commercial or regulatory compliance, any specimen tested as directed in the monograph for that article shall comply.

Tests and assays in this Pharmacopeia prescribe operation on a single specimen, that is, the singlet determination, which is the minimum sample on which the attributes of a compendial article should be measured. Some tests, such as those for *Dissolution* and *Uniformity of dosage units*, require multiple dosage units in conjunction with a decision scheme. These tests, albeit using a number of dosage units, are in fact the singlet determinations of those particular attributes of the specimen. These procedures should not be confused with statistical sampling plans. Repeats, replicates, statistical rejection of outliers, or extrapolations of results to larger populations are neither specified nor proscribed by the compendia; such decisions are dependent on the objectives of the testing. Commercial or regulatory compliance testing, or manufacturer's release testing, may or may not require examination of additional specimens, in accordance with predetermined guidelines or sampling strategies. Treatments of data handling are available from organizations such as ISO, IUPAC, and AOAC.

Where the *Content Uniformity* determinations have been made using the same procedure specified in the *Assay*, the average of all of the individual *Content Uniformity* determinations may be used as the *Assay* value.

**Description**—Information on the "description" pertaining to an article, which is relatively general in nature, is provided in the reference table *Description and Relative Solubility of USP and NF Articles* in this Pharmacopeia for those who use, prepare, and dispense drugs and/or related articles, solely to indicate properties of an article complying with monograph standards. The properties are not in themselves standards or tests for purity even though they may indirectly assist in the preliminary evaluation of an article.

**Solubility**—The statements concerning solubilities given in the reference table *Description and Relative Solubility of USP and NF Articles* for Pharmacopeial articles are not standards or tests for purity but are provided primarily as information for those who use, prepare, and dispense drugs and/or related articles. Only where a quantitative solubility test is given, and is designated as such, is it a test for purity.

The approximate solubilities of Pharmacopeial substances are indicated by the descriptive terms in the accompanying table.

Descriptive Term	Parts of Solvent Required for 1 Part of Solute
Very soluble	Less than 1
Freely soluble	From 1 to 10
Soluble	From 10 to 30
Sparingly soluble	From 30 to 100
Slightly soluble	From 100 to 1000
Very slightly soluble	From 1000 to 10,000
Practically insoluble, or Insoluble	Greater than or equal to 10,000

Soluble Pharmacopeial articles, when brought into solution, may show traces of physical impurities, such as minute fragments of filter paper, fibers, and other particulate matter, unless limited or excluded by definite tests or other specifications in the individual monographs.

**Interchangeable Methods**—Certain general chapters contain a statement that the text in question is harmonized with the corresponding text of the *European Pharmacopoeia* and/or the *Japanese Pharmacopoeia* and that these texts are interchangeable. Therefore, if a substance or preparation is found to comply with a requirement using an interchangeable method from one of these pharmacopeias, it should comply with the requirements of the *United States Pharmacopeia*. However, where a difference appears, or in the event of dispute, only the result obtained by the procedure given in this Pharmacopeia is conclusive.

## PREScribing AND DISPENSING

Prescriptions for compendial articles shall be written to state the quantity and/or strength desired in metric units unless otherwise indicated in the individual monograph (see also *Units of Potency* in these *General Notices*). If an amount is prescribed by any other system of measurement, only an amount that is the metric equivalent of the prescribed amount shall be dispensed.

## PRESERVATION, PACKAGING, STORAGE, AND LABELING

**Containers**—The *container* is that which holds the article and is or may be in direct contact with the article. The *immediate container* is that which is in direct contact with the article at all times. The *closure* is a part of the container.

Prior to being filled, the container should be clean. Special precautions and cleaning procedures may be necessary to ensure that each container is clean and that extraneous matter is not introduced into or onto the article.

The container does not interact physically or chemically with the article placed in it so as to alter the strength, quality, or purity of the article beyond the official requirements.

The Pharmacopeial requirements for the use of specified containers apply also to articles as packaged by the pharmacist or other dispenser, unless otherwise indicated in the individual monograph.

**Tamper-Evident Packaging**—The container or individual carton of a sterile article intended for ophthalmic or otic use, except where extemporaneously compounded for immediate dispensing on prescription, shall be so sealed that the contents cannot be used without obvious destruction of the seal.

Articles intended for sale without prescription are also required to comply with the tamper-evident packaging and labeling requirements of the FDA where applicable.

Preferably, the immediate container and/or the outer container or protective packaging used by a manufacturer or distributor for all dosage forms that are not specifically exempt is designed so as to show evidence of any tampering with the contents.

**Light-Resistant Container** (see *Light Transmission* under *Containers* (661))—A light-resistant container protects the contents from the effects of light by virtue of the specific properties of the material of which it is composed, including any coating applied to it. Alternatively, a clear and colorless or a translucent container may be made light-resistant by means of an opaque covering, in which case